



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(51) International Patent Classification ⁶ : C07K 14/715		A2	(11) International Publication Number: WO 96/17869
			(43) International Publication Date: 13 June 1996 (13.06.96)
(21) International Application Number: PCT/IT95/00208		(81) Designated States: AU, CA, CN, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 5 December 1995 (05.12.95)			
(30) Priority Data: RM94A000794 6 December 1994 (06.12.94) IT		Published Without international search report and to be republished upon receipt of that report.	
(71) Applicant (for all designated States except US): ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. ANGELETTI S.P.A. [IT/IT]; Via Pontina Km. 30.600, I-00040 Pomezia (IT).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): CILIBERTO, Gennaro [IT/IT]; Viale Gorgia di Leontini, 330/19, I-00124 Casalpalocco (IT). TONIATTI, Carlo [IT/IT]; Via Benedetto Croce, 26, I-00142 Roma (IT).			
(74) Agents: DI CERBO, Mario et al.; Società Italiana Brevetti S.p.A., Piazza di Pietra, 39, I-00186 Roma (IT).			

(54) Title: INTERLEUKIN-6 (IL-6) ANTAGONISTS

(57) Abstract

These are antagonists of interleukin-6, characterised in that they consist of soluble forms of the receptor α of human IL-6 (shIL-6R α) containing one or more mutation in the gp130 binding interface. In a preferred embodiment, the mutations are present in a position chosen from the group comprising Ala 228, Asn 230, His 280 and Asp 281. These antagonists can be used as agents capable of preventing and treating diseases caused by abnormal IL-6 activity. The figure shows the antagonist activity of the mutant Ala228Asp/Asn 230Asp/His280Ser/Asp281Val in comparison with the agonist properties of wild type shIL-6R α . This antagonist can be used for preparation of drugs for the prevention, control and treatment of diseases caused by abnormal IL-6 bioactivity.

	Wild type					SEQ ID NO:4				
shIL-6R α (100 nM)	—	—	—	—	+	+	+	+	+	+
hIL-6 (pM)	—	2	20	200	2	20	200	2	20	200



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INTERLEUKIN-6 (IL-6) ANTAGONISTS

DESCRIPTION

5 The present invention relates to the field of immunology. More specifically, the subject of the present invention are mutants of the soluble forms of receptor α of interleukin 6 that interfere in a negative manner in the formation of the dimeric receptor complex
10 between the monomeric receptor complex IL-6/sIL-6R α and gp 130. A further subject of the invention is the use of these mutants as interleukin-6 antagonists to control, prevent and treat the diseases caused by abnormal IL-6 activity.

15 The term "interleukin 6" or "IL-6", in the context of the present invention, means IL-6 and fragments, deletions, insertions, substitutions, mutations and modifications thereof that maintain the biological characteristics of wild IL-6. Unless otherwise
20 specified, the term refers to human IL-6.

 Interleukin-6 elicits a variety of biological responses on different target cells. However - while the physiological production of IL-6 regulates B-cells proliferation and maturation, T-cell activation and the
25 production of acute-phase proteins in liver during inflammatory response - dysregulated production of the cytokine plays a crucial role in the pathogenesis of many inflammatory, autoimmune and neoplastic diseases.

 As is known from the state of the art, various
30 attempts have been made to perfect IL-6 bioactivity inhibitors, both in order to study the role played by this cytokine in the genesis of single diseases, and in order to define drugs for treatment.

 It is also known that the mature human IL-6 protein
35 (h IL-6) is a 185 amino acid polypeptide containing two disulfide bonds (Cys 45 - Cys 51, and Cys 74 - Cys 84). IL-6 functions through interaction of two binding sites

(known as site I and site II) with at least two specific receptors (IL-R α and gp 130) on the surface of the target cells, creating a trimeric complex IL-6-(receptor)₂. This complex is formed sequentially: a first receptor
5 (IL-6R α) binds with low affinity to site I of IL-6 without transmitting the signal; subsequently a second receptor (gp 130), after binding with high affinity to site II of IL-6, transduces the signal.

Based on this mechanism, hIL-6 mutants have been
10 designed that are capable of binding a first receptor to site I, but incapable of dimerising the receptor because of mutations that sterically inhibit binding of the second receptor to site II. Mutants of this type have been described in WO 94/09138 (Cetus Oncology Corporation), and WO 94/011402 and PCT/IT 94/00095
15 (Istituto di Ricerche di Biologia Molecolare P. Angeletti S.p.A.).

It has now surprisingly been found, and this discovery forms the basis of the present invention, that
20 soluble forms of the receptor α of IL-6 (sIL-6R α) containing one or more mutations in the region interfacing with gp 130 are antagonists of interleukin 6.

In one embodiment, sIL-6R α contains at least one mutation in a position chosen from Ala 228, Asn 230, His
25 280 and Asp 281. The mutations may, for example, be chosen from the group comprising: Asn230Asp (SEQ ID NO:1); Ala228Asp/Asn230Asp (SEQ ID NO:2); His280Ser/Asp281Val (SEQ ID NO:3).

In a preferred embodiment, sIL-6R α contains multiple
30 mutations in positions Ala 228, Asn 230, His 280 and Asp 281. A multiple mutation that has given good results is Ala228Asp/Asn230Asp/His280Ser/Asp281Val (SEQ ID NO:4).

The antagonists of IL-6, which according to the invention consist of soluble forms of the receptor α of
35 IL-6, mutated in the interface that binds to gp 130, can be administered at concentrations therapeutically effective for the treatment and prevention of diseases

related to abnormal IL-6 activity. For this purpose, the antagonists according to the invention are administered preferably by endovenous and/or subcutaneous injection. Experts in the field are well acquainted with these methods of administration.

Up to this point a general description of the invention has been given. A more detailed description of the invention will now be provided, with the aid of the following examples, in order to give a clearer understanding of the objects, characteristics, advantages and operating methods thereof.

Figure 1A shows resolution on SDS/PAGE gel (12%), in vitro, of the co-immunoprecipitates of mutants according to the invention with sgp 130 (soluble form of gp 130) marked using ^{35}S . Figure 1B shows the position on the gel corresponding to the migrated ^{125}I -IL-6.

Figure 2 shows, in the form of a histogram, the different capacities of the wild type sIL-6R α receptor and the sIL-6R α receptors mutated to interact with gp 130 on the membrane of A 375 cells.

Figure 3 shows resolution on SDS/PAGE gel (12%), in vitro, of the co-immunoprecipitates of sgp 130 marked with ^{35}S or with 2.0 μg of wild receptor or with 2.0 μg of mutant receptor (SEQ ID NO: 4).

Figure 4 shows the antagonist activity of the mutant SEQ ID NO: 4 on HepG2 cells in comparison with the agonistic properties of wild type shIL-6R α .

Figure 5 shows resolution on gel of the complex formed by APRF with the DNA binding site (SIE, Serum Inducible Element). (The mutant SEQ ID NO: 4 does not inhibit activation of OM-dependent APRF on HepG2 cells).

Antagonists of IL-6 according to the invention may be produced synthetically or using recombinant techniques. In the latter case the cDNA coding for sIL-6R α can be incorporated in a plasmid before being expressed in procaryotic or eukaryotic cells. Bacteria are the preferred procaryotic microorganisms.

Alternatively, the cDNA coding for the sIL-6R α mutants according to the invention can be introduced into mammal cells: these mammal cells can be chosen from the group comprising CHO, COS, C127, HepG2, SK Hep.
5 Furthermore, the protein can also be expressed in insect cells (Sf9 or HighFive) using the well known recombinant Baculovirus technology (AcNPV strain).

Example 1

Preparation of sIL-6R α mutants

10 sIL-6R α cDNA was obtained as a 5'EcoRI-3'XbaI fragment by PCR using the IL-6R α complete cDNA as template (Sporeno E., Paonessa G., Salvati A.L., Graziani R., Delmastro P., Ciliberto G. and Toniatti C. (1994) J. Biol. Chem. 269, 10991-10995). The 3' primer was
15 designed in order to introduce an artificial TAG stop codon at amino acid position 324 preceded by a six-histidine coding sequence. This was done to ensure that the sIL-6R α and the mutants thereof produced by us have a six histidine "tail" at the carboxy-terminal end of the
20 molecule, which can be useful for purification of the molecule using metal affinity chromatography. The fragment generated was then introduced into the COS-7 cells expression vector pcDNA1 (Invitrogen) and sequenced in its entirety, thus obtaining plasmid pC6FRH.

25 Plasmid pC6FRH was in turn used as a template to obtain constructs coding for the following four mutants: Asn230Asp (SEQ ID NO: 1); Asn230Asp/Ala228Asp (SEQ ID NO: 2); His280Ser/Asp281Val (SEQ ID NO 3) and
30 Ala228Asp/Asn230Asp/His280Ser/Asp281Val (SEQ ID NO: 4), by a two-step PCR as described in the past (Landt O., Grunert H.P., and Hahn, U. (1990) Gene 96, 125-128).

The mutants were then expressed in COS-7 cells. For this purpose, the COS-7 cells were maintained in Dulbecco's modified Eagle's medium supplemented with 10%
35 FCS, plus glutamine and antibiotics, at 5% CO₂. For protein expression, 2.5x10⁶ COS-7 cells were seeded in 100-mm tissue culture dishes and the next day transfected

with 2 μ g of the various shIL-6R α expression vectors using the DEAE-dextran technique as described in Seed B., Aruffo A. (1987) Proc. Natl. Acad. Sci. USA 84, 3365-3369. 16 h after transfection cells were split, replated
5 in 100-mm dishes and grown in complete medium at 37°C. After 72-96 h the medium was collected, centrifuged and used for co-immunoprecipitation experiments and binding analysis. To monitor the expression level of each mutant, 2.5x10⁵ transfected COS-7 cells were replated in
10 35-mm dishes and, 48 h after transfection, metabolically labeled with [³⁵S]methionine for 4h. The supernatants were immunoprecipitated with anti-human IL-6R α monoclonal antibody I6R1/9.G11. The immunoprecipitated material was then analyzed by electrophoresis through a polyacrylamide
15 gel containing SDS.

IL-6 binding analysis of mutants

To determine the affinity for IL-6 of the sIL-6R α mutants contained in the conditioned COS cells culture medium, the six-histidine "tail" in the proteins produced
20 by us was used.

Appropriate amounts of transfected COS cells supernatant, previously determined in titration experiments and to which imidazole was added up to a final concentration of 5 mM, were mixed with 20-40 pM
25 ¹²⁵I-IL-6 and increasing concentrations of unlabeled cytokine. At equilibrium conditions, 40 μ l of Ni²⁺-NTA-Agarose (Quiagen), a resin capable of selectively binding proteins containing histidine "tails", was added and incubation prolonged for one additional hour. The
30 ligand bound by the receptor and, therefore, indirectly resin-associated, was separated from free ligand (supernatant) by centrifugation through a cushion of 30% sucrose in PBS. All steps were performed at 4°C. The values of free and bound cpm (radioactivity count per
35 minute) allowed drawing of competition displacement curves. The apparent K_d of the various soluble IL-6 receptors was determined after Scatchard transformation

of the results (see the article by Sporeno, Paonessa, Salviati, Graziani, Delmastro, Ciliberto and Toniatti (1994), J. Biol. Chem. 269, 10991-10995). Analysis of binding data and curve fitting was done using the
5 UltraFit software (Biosoft®) on a Macintosh computer.

As can be seen from Table 1, the binding affinity:
1) remains substantially unprejudiced for mutants Asn230Asp (SEQ ID NO: 1), Asn230Asp/Ala228Asp (SEQ ID NO: 2); b) is reduced in a slight but reproducible manner
10 (between 2.5 nM and 6 nM) for His280Ser/Asp281Val (SEQ ID NO: 3); and c) the activity of the wild type form is fully maintained for the mutant Ala228Asp/Asn230Asp/His280Ser/Asp281Val (SEQ ID NO: 4).

15 TABLE 1
IL-6 binding affinity of soluble interleukin 6 receptor
(wild and mutated), labeled with histidine and expressed
in COS cells

Mutant	IL-6 binding affinity
	nM
Asn230Asp	(SEQ ID NO: 1) 1.5 ± 1
Ala228Asp/Asn230Asp	(SEQ ID NO: 2) 2.0 ± 1
His280Ser/Asp281Val	(SEQ ID NO: 3) 4.3 ± 2
25 Ala228Asp/Asn230Asp/ His280Ser/Asp281Val	(SEQ ID NO: 4) 2.5 ± 1
wild type sIL-6Rα	2.0 ± 1

Binding of mutants to gp130

The IL-6-dependent formation of sIL-6Rα/sgp130
30 heterodimers can be easily monitored in vitro by co-immunoprecipitations with the use of suitable monoclonal antibodies. A selection was therefore made of the mutants showing an IL-6 binding affinity in the same order of magnitude as the wild-type and their binding to
35 sgp130 was evaluated by co-immunoprecipitation in the presence of IL-6.

³⁵S-labelled sgp 130 was incubated with ¹²⁵I-IL-6 (as an internal standard for the immunoprecipitation) and aliquots of transfected COS-7 cell culture medium, containing either native or mutant receptors. After in vitro binding, anti IL-6Rα mAb I6/R19.G11 was added to the mixtures and the immunoprecipitates resolved on SDS/PAGE gels. The results are shown in figure 1. As can be seen, the mutant His280Ser/Asp281Val (SEQ ID NO: 3) strongly reduced the association of the IL-6/ sIL-6Rα complex sgp 130 molecule (lane 4). Also the single substitution mutant Asn230Asp (SEQ ID NO: 1) had decreased interaction with sgp130 (lane 2). The effect of this mutant on interaction with sgp 130 is of the same magnitude as that of mutant Ala228Asp/Asn230Asp (SEQ ID NO: 2). Finally, the quadruple mutant Ala228Asp/Asn230Asp/His280Ser/Asp281Val (SEQ ID NO: 4) displayed the lowest ability to co-immunoprecipitate ³⁵S-gp 130 (lane 5).

The receptor mutants were also tested for their ability to interact with gp130 on the cell surface. For this purpose, binding experiments were performed on human melanoma A375 cells which have an excess of gp130 molecules over IL-6Rα. In fact, binding of ¹²⁵I-IL-6 to A375 monolayers could be strongly enhanced by addition of the soluble receptor. This phenomenon is due to the ability of the sIL-6Rα to bind ¹²⁵I-IL-6 and subsequently to interact with the gp130 molecules present on the surface of the cells. The specificity of this interaction is demonstrated by the fact that the increased binding of ¹²⁵I-IL-6 in the presence of sIL-6Rα is competed by the addition of an excess of unlabeled human Oncostatin M (OM) (fig. 2). This cytokine is able to bind directly to gp130 and to compete the binding of the IL-6/IL-6Rα complex with gp130. Unlike wild type shIL-6Rα, only a minor increase in specific binding was detected when cells were challenged with ¹²⁵I-IL-6 plus the soluble mutant receptors acting as antagonists

(figure 2). The greatest disturbance to interaction with gp130 is shown by the mutants His280Ser/Asp281Val (SEQ ID NO: 3) and Ala228Asp/Asn230Asp/His280Ser/Asp281Val (SEQ ID NO: 4).

5 Demonstration that the mutant Ala228Asp/Asn230Asp/His280Ser/Asp281Val (SEQ ID NO: 4) is an IL-6 antagonist

10 The results shown above indicate this mutant to be the one that, without any decrease in the affinity for IL-6, has the greatest effect on gp130 binding. In order to study the biological activity of this mutant, its production and that of the wild type soluble receptor was scaled up using the MaxBac® system (Invitrogen's Baculovirus Expression System). The expressed and purified receptors were tested in immunoprecipitation
15 experiments.

20 Sf9 cells, grown in Grace's insect medium, were used for transfection of transfer vectors, isolation of recombinant virus and preparation of high-titer virus stocks. High Five® cells (Invitrogen) were instead used for production of proteins. Briefly, 4×10^7 High Five® cells, grown in complete Grace's insect medium, were seeded into 750 ml flasks and infected with the appropriate recombinant virus at a MOI of 10. After 2 hours, cells were washed and SF-900 serum-free medium was
25 added. The culture supernatants were harvested at 36 hours post-infection, dialyzed against PBS and directly loaded on a Ni^{2+} -NTA-agarose column. After washing steps with PBS/8 mM imidazole, both wild type shIL-6R α and mutant were eluted in PBS/80 mM imidazole. Purified
30 protein was dialyzed against PBS and directly used or stored at 4°C for up to three weeks.

35 The use of purified proteins made it possible to quantify the amount of receptor and to compare the ability of wild type receptor and mutant receptor to interact with sgpl30 in the presence of a wide range of IL-6 concentrations. The results (figure 3) were in good agreement with those previously obtained with COS cell-

derived receptors. Interestingly at each point of the curve the amount of co-immunoprecipitated gp130 is lower for the mutant receptor than for the wild type receptor and this is true even at the highest concentrations of the cytokine tested (100 nM). These findings confirm that the interaction between IL-6 and mutant sIL-6R α generates a complex with lower affinity for gp130.

To test the potential antagonism of the mutant for IL-6 activity, we chose the IL-6 dependent activation of transcription factor APRF (Acute Phase Response Factor)/STAT 3 in human hepatoma cells. Activation of APRF, which is dependent on tyrosine phosphorylation, is a rapid cytoplasmic event which takes place within minutes of stimulation with all cytokines of the IL-6 family.

Human HepG2 cells were stimulated with increasing concentrations of IL-6 together with a fixed amount of wild type and mutant receptors (100 nM). As shown in figure 4, while shIL-6R α potentiated APRF induction by IL-6 (compare lanes 2-4 and 5-7 in figure 4), the soluble mutant receptor not only lacked any agonistic activity but instead down-modulated the activity of the cytokine (figure 4, lanes 8-10). As predicted from the co-immunoprecipitation experiments, the inhibition was complete when cells were treated with up to 20 pM IL-6 (lanes 8 and 9, figure 4) and only a weak induction was detected at the highest concentration of IL-6 (200 pM, lane 10 of figure 4).

To test if antagonism was specific for IL-6, the mutant was added to HepG2 cells induced with the IL-6-related cytokine OM, which is also known to efficiently induce APRF phosphorylation. As shown in figure 5, the mutant did not antagonize OM activity.

SEQUENCE LISTING

GENERAL INFORMATION

- (i) APPLICANT: ISTITUTO DI RICERCHE DI BIOLOGIA
MOLECOLARE P. ANGELETTI S.p.A.
- 5 (ii) TITLE OF INVENTION:
(iii) NUMBER OF SEQUENCES:
(iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Società Italiana Brevetti
(B) STREET: Piazza di Pietra, 39
10 (C) CITY: Rome
(D) COUNTRY: Italy
(E) POSTAL CODE: I-00186
(v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk 3.5" 1.44 MBYTES
15 (B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS Rev. 6.22
(D) SOFTWARE: Microsoft Word 6.0
(viii) ATTORNEY INFORMATION
(A) NAME: DI CERBO, Mario (Dr.)-
20 (C) REFERENCE: RM/X88470/PC-DC
(ix) TELECOMMUNICATION INFORMATION
(A) TELEPHONE: 06/6785941
(B) TELEFAX: 06/6794692
(C) TELEX: 612287 ROPAT
25 (1) INFORMATION FOR SEQ ID NO: 1:
(i) SEQUENCE CHARACTERISTICS
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
30 (ii) MOLECULE TYPE: protein
(iii) HYPOTHETICAL: no
(v) FRAGMENT TYPE: internal
(vii) IMMEDIATE SOURCE:
(A) SYNTHESIS: production as recombinant
35 protein in eucariotic cells
(ix) FEATURE:
(A) NAME: Asn230Asp

- 11 -

(C) IDENTIFICATION

METHOD:

Electrophoresis on a denaturing SDS/polyacrylamide gel

(D) OTHER INFORMATION: amino acid

sequence of a mutant form of the interleukin 6 receptor
 5 from position 222 to position 236.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Ile Thr Val Thr Ala Val Ala Arg Asp Pro Arg Trp Leu Ser Val
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO: 2:

10 (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

15 (iii) HYPOTHETICAL: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:

(A) SYNTHESIS: production as recombinant
 protein in eucariotic cells

20 (ix) FEATURE:

(A) NAME: Ala228Asp/Asn230Asp

(C) IDENTIFICATION

METHOD:

Electrophoresis on a denaturing SDS/polyacrylamide gel

(D) OTHER INFORMATION: amino acid

25 sequence of a mutant form of the interleukin 6 receptor
 from position 222 to position 236.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Ile Thr Val Thr Ala Val Asp Arg Asp Pro Arg Trp Leu Ser Val
 1 5 10 15

30 (3) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: no

(v) FRAGMENT TYPE: internal

- 12 -

- (vii) IMMEDIATE SOURCE:
 (A) SYNTHESIS: production as recombinant protein in eucariotic cells
- (ix) FEATURE:
 5 (A) NAME: His280Ser/Asp281Val
 (C) IDENTIFICATION METHOD: Electrophoresis on a denaturing SDS/polyacrylamide gel
 (D) OTHER INFORMATION: amino acid sequence of a mutant form of the interleukin 6 receptor
 10 from position 273 to position 287.
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:
 Leu Gln His His Cys Val Ile Ser Val Ala Trp Ser Gly Leu Arg
 1 5 10 15
- (4) INFORMATION FOR SEQ ID NO: 4:
 15 (i) SEQUENCE CHARACTERISTICS
 (A) LENGTH: 66 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein
 20 (iii) HYPOTHETICAL: no
 (v) FRAGMENT TYPE: internal
 (vii) IMMEDIATE SOURCE:
 (A) SYNTHESIS: production as recombinant protein in eucariotic cells
- (ix) FEATURE:
 25 (A) NAME:
 Ala228Asp/Asn230Asp/His280Ser/Asp281Val
 (C) IDENTIFICATION METHOD: Electrophoresis on a denaturing SDS/polyacrylamide gel
 30 (D) OTHER INFORMATION: amino acid sequence of a mutant form of the interleukin 6 receptor from position 222 to position 287.
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:
 Ile Thr Val Thr Ala Val Asp Arg Asp Pro Arg Trp Leu Ser Val Thr
 35 1 5 10 15
 Trp Gln Asp Pro His Ser Trp Asn Ser Ser Phe Tyr Arg Leu Arg Phe
 20 25 30

- 13 -


	Glu	Leu	Arg	Tyr	Arg	Ala	Glu	Arg	Ser	Lys	Thr	Phe	Thr	Thr	Trp	Met
			35					40					45			
	Val	Lys	Asp	Leu	Gln	His	His	Cys	Val	Ile	Ser	Val	Ala	Trp	Ser	Gly
		50					55					60				
5	Leu	Arg														
	65															

CLAIMS

1. Antagonist of interleukin-6 (IL-6), characterized in that it consists in soluble forms of the α receptor of IL-6 (sIL-6R α) containing one or more mutations in the interface binding with gp 130.
5
2. Antagonist of interleukin-6 according to claim 1, in which sIL-6R α contains at least one mutation in a position chosen from the group comprising Ala 228, Asn 230, His 280 and Asp 281.
- 10 3. Antagonist of interleukin 6 according to claim 1 or 2, in which sIL-6R α contains the mutation Asn230Asp (SEQ ID NO: 1).
4. Antagonist of interleukin 6 according to claim 1 or 2, in which sIL-6R α contains the mutation
15 Ala228Asp/Asn230/Asp (SEQ ID NO: 2).
5. Antagonist of interleukin 6 according to claim 1 or 2, in which sIL-6R α contains the mutations His280Ser/Asp281Val (SEQ ID NO: 3).
6. Antagonist of interleukin 6 according to claim 1
20 or 2, in which sIL-6R α contains the mutations Ala228Asp/Asn230Asp/His280Ser/Asp281Val (SEQ ID NO: 4).
7. Use of interleukin 6 antagonists according to claims 1 to 6, for the study and preparation of drugs capable of controlling, preventing and treating diseases
25 caused by abnormal IL-6 activity.
8. Soluble receptors acting as IL-6 antagonists and their use, as described, illustrated and claimed above.


1/5

Fig. 1




2/5

FIG. 2



3/5

Fig. 3



4/5

Fig. 4



	Wild type						SEQ ID NO:4			
shIL-6R α (100 nM)	—	—	—	—	+	+	+	+	+	+
hIL-6 (pM)	—	2	20	200	2	20	200	2	20	200



1 2 3 4 5 6 7 8 9 10

5/5

Fig. 5





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/715		A3	(11) International Publication Number: WO 96/17869
			(43) International Publication Date: 13 June 1996 (13.06.96)
(21) International Application Number: PCT/IT95/00208		(81) Designated States: AU, CA, CN, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 5 December 1995 (05.12.95)			
(30) Priority Data: RM94A000794 6 December 1994 (06.12.94) IT		Published With international search report.	
(71) Applicant (for all designated States except US): ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. ANGELETTI S.P.A. [IT/IT]; Via Pontina Km. 30.600, I-00040 Pomezia (IT).		(88) Date of publication of the international search report: 29 August 1996 (29.08.96)	
(72) Inventors; and (75) Inventors/Applicants (for US only): CILIBERTO, Gennaro [IT/IT]; Viale Gorgia di Leontini, 330/19, I-00124 Casalpalocco (IT). TONIATTI, Carlo [IT/IT]; Via Benedetto Croce, 26, I-00142 Roma (IT).			
(74) Agents: DI CERBO, Mario et al.; Società Italiana Brevetti S.p.A., Piazza di Pietra, 39, I-00186 Roma (IT).			

(54) Title: INTERLEUKIN-6 (IL-6) ANTAGONISTS

(57) Abstract

These are antagonists of interleukin-6, characterised in that they consist of soluble forms of the receptor α of human IL-6 (shIL-6R α) containing one or more mutation in the gp130 binding interface. In a preferred embodiment, the mutations are present in a position chosen from the group comprising Ala 228, Asn 230, His 280 and Asp 281. These antagonists can be used as agents capable of preventing and treating diseases caused by abnormal IL-6 activity. The figure shows the antagonist activity of the mutant Ala228Asp/Asn 230Asp/His280Ser/Asp281Val in comparison with the agonist properties of wild type shIL-6R α . This antagonist can be used for preparation of drugs for the prevention, control and treatment of diseases caused by abnormal IL-6 bioactivity.

	Wild type					SEQ ID NO:4				
shIL-6R α (100 nM)	—	—	—	—	+	+	+	+	+	+
hIL-6 (pM)	—	2	20	200	2	20	200	2	20	200



1 2 3 4 5 6 7 8 9 10

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INTERNATIONAL SEARCH REPORT

International Application No
PLI/IT 95/00208A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K14/715

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBO J., vol. 12, no. 4, 1993, pages 1705-1712, XP002003859 YAWATA ET AL.: "Structure-function analysis of human IL-6 receptor : dissociation of amino acid residues required for IL-6-binding and IL-6 signal transduction through gp130" see page 1708 - page 1711 --- -/-	1,2,5,7, 8

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

24 May 1996

Date of mailing of the international search report

07.06.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Gac, G

INTERNATIONAL SEARCH REPORT

International Application No
PC/IT 95/00208

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Week 9320 Derwent Publications Ltd., London, GB; AN 93-161739 XP002003861 & JP,A,05 091 892 (CHUGAI PHARM CO LTD & AL) , 16 April 1993 see abstract & DATABASE STRAND TPSD AN: E52907, see in "mutation" list. ---	1,2,5,7
P,X	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 20, 19 May 1995, pages 12242-12249, XP000566420 SALVATI ET AL.: "Interleukin-6 (IL-6) antagonism by soluble IL-6 receptor alpha mutated in the predicted gp130-binding interface" see the whole document ---	1-8
A	COLD SPRING HARBOR SYMPOSIA ON QUANTITATIVE BIOLOGY, vol. 54, 1989, pages 713-722, XP002003860 TAGA ET AL.: "Interleukin-6 receptor and a unique mechanism of its signal transduction" see the whole document ---	1,7,8
A	EUR. CYTOKINE NETWORK, vol. 5, no. 3, May 1994 - June 1994, pages 293-300, XP000571448 LIAUTARD ET AL.: "Epitope analysis of human IL-6 receptor gp80 molecule with monoclonal antibodies" see the whole document -----	1,7,8